

The chart is orientated for
Aug. 15 at 10 p.m.
Sep. 1 at 9 p.m.
Sep. 15 at 8 p.m.

Evening sky in September 2025

To use the chart, hold it up to the sky. Turn the chart so the direction you are looking is at the bottom of the chart. If you are looking to the south then have 'South horizon' at the lower edge. As the earth turns the sky appears to rotate clockwise around the south celestial pole (SCP on the chart). Stars rise in the east and set in the west, just like the sun. The sky makes a small extra clockwise rotation each night as we orbit the sun.

Canopus twinkles like a diamond near the southern horizon. Vega shines on the northern horizon. Orange Arcturus twinkles red and green as it sets in the northwest. The Milky Way spans the sky from north to south. Mars is a medium-bright red 'star' low in the west. Saturn is a cream-coloured 'star' low in the east. West of overhead is orange Antares marking the Scorpion's body. The Scorpion's tail curls above it like a back-to-front question mark. Crux, the Southern Cross, and the Pointers are in the south-west. There are eclipses of both the Moon and the Sun this month.

The Evening Sky in September 2025

Bright stars shine around the skyline. Orange **Arcturus** is in the northwest, often twinkling red and green as it sets. **Canopus**, the brightest star in the evening sky, skims along the southern skyline, twinkling all colours. **Vega** is low in the north. It is the second-brightest northern star after Arcturus. From northern Aotearoa the star **Deneb** can be seen near the north skyline. It is the brightest star in Cygnus the Swan.

Mars is low in the west. It looks like a medium-bright red star. In mid-September it will be near **Spica**, the brightest star in Virgo. **Saturn** is in the eastern sky, a medium-bright 'star' all on its own. Late in the month Mercury appears below Mars in the west, setting an hour after the Sun (so isn't on the chart.)

Orange **Antares**, northwest of the zenith, marks the body of the Scorpion. The Scorpion's tail hooks toward the zenith like a back-to-front question mark. It is the 'fishhook of Maui' in Māori star lore. Below or right of the Scorpion's tail is 'the teapot' made by the brightest stars of **Sagittarius**. It is upside down in our southern hemisphere view.

Midway down the southwest sky are 'The Pointers', Beta and **Alpha Centauri**. They point down to **Crux** the Southern Cross. Alpha Centauri is the third brightest star. It is also the closest of the naked-eye stars, 4.3 light-years* away. Beta Centauri, along with most of the stars in Crux, is a blue-giant star hundreds of light-years away.

The **Milky Way** spans the sky from north to south. It is brightest and broadest overhead in Scorpius and Sagittarius. In a dark sky it can be traced down past the Pointers and Crux into the southwest. To the northeast it passes **Altair**, meeting the skyline right of **Vega**. The Milky Way is our edgewise view of the galaxy, the pancake of billions of stars of which the sun is just one. The thick hub of the galaxy, 27 000 light years away, is in Sagittarius. Dust clouds near us appear as gaps and slots in the Milky Way. Binoculars show many clusters of stars and some glowing gas clouds in the Milky Way.

The Large and Small Clouds of Magellan, **LMC** and **SMC**, look like two misty patches of light in the south sky. They are easily seen by eye on a dark moonless night. They are galaxies like our Milky Way but much smaller. The LMC is about 160 000 light years away; the SMC about 200 000 light years away.

On moonless evenings in a dark sky the Zodiacal Light is visible in the west. It appears as a faint broad column of light extending up past Mars and Spica, toward Antares. It is sunlight reflecting off meteoric dust in the plane of the solar system.

There are two 'morning stars'. Golden Jupiter appears in the northeast around 4:20 a.m. at the beginning of the month and 3 a.m. at the end. Silver Venus, brighter than Jupiter, rises around 5:30 at the beginning of September and soon after 5 a.m. at the end. The Moon will be near Jupiter on the 14th, and close to Venus on the 20th.

There is a **total eclipse of the Moon** around dawn on Monday 8th. The Moon starts to enter the dark part of Earth's shadow, the umbra, at 4:27 a.m. It is fully in the umbra at 5:31 and closest to the shadow centre at 6:13. It starts moving out of the umbra at 6:53, around moonset from N.Z.

A **partial eclipse of the Sun** happens at sunrise on Monday 22nd. There will be a big bite out of the Sun when it rises. This will get bigger till 7:10 a.m., then slowly shrink. The Moon moves off the Sun around 8:15, depending on your location. **Never look at the Sun without eye protection!** Doing so can cause permanent damage. Eclipse glasses are the best protection. If they aren't available then make pinhole in a sheet of paper or cardboard and project an image of the sun onto a wall. Eclipse glasses can be bought from Astronz, Auckland, www.astronz.nz; Carter Observatory = Space Place, Wellington <https://shopwellington.nz/products/0a6f6e36-8b23-11eb-f3d6-6a8b0d828055>; the Dark Sky Project, Lake Tekapo, www.darkskyproject.co.nz, and Otago Museum, Samanta.Luzzi@tuhura.nz .

*A **light-year (l.y.)** is the distance that light travels in one year: nearly 10 million million km or 10^{13} km. Sunlight takes eight minutes to get here; moonlight about one second. Sunlight reaches Neptune, the outermost major planet, in four hours. It takes sunlight four years to reach the nearest star, Alpha Centauri.



Interesting Objects West of Overhead in September

Antares is the brightest star in the region. It is orange coloured; being a 'red giant' star. (The 'red' of red giants is usually more an orange tint.) It is around 600 light years* away, 19 000 times brighter than the sun, and three times bigger than Earth's orbit. Its mass or weight is about 12 times that of the sun, so most of the star is very thin gas spread around a hot dense core. Red giants are the last stage in the evolution of stars. The dense core of the star has shrunk and heated. The outer regions of the star have expanded to a very spread-out gas. The core is wringing the last of the thermo-nuclear energy out of elements like helium, carbon, oxygen and neon. Relatively soon the core of Antares will run out of energy and collapse, triggering a spectacular supernova explosion. (The sun will become a red-giant in about seven billion years time but it ends up as a white dwarf star, not a supernova.)




At the right-angle bend in the Scorpion's tail is a large and bright cluster of stars, **NGC 6231**, looking like a small comet. It is around 6000 l.y. away. Its brightest stars are 60 000 times brighter than the sun. The cluster is about 8 light years across, similar in size to the Pleiades/Matariki cluster in our summer sky. Were it as close as the Pleiades/Matariki cluster (440 l.y.) then its brightest stars would be as bright as Sirius.

Below and right of the Scorpion's sting is M7, a cluster obvious to the eye and nicely seen in binoculars. M7 is about 800 l.y. away and around 260 million years old. Below the sting and fainter is M6, the 'butterfly cluster'. M6 is around 1300 l.y. away and is half the age of M7. Other clusters worth a look in binoculars are M21, M23, NGC 6167, and NGC 6193. The 'M' objects were listed by the 18th Century French astronomer Charles Messier. He hunted comets, so made a catalogue of fuzzy objects that could be mistaken for comets. The NGC (New General Catalogue) objects shown are bright enough to have been seen by Messier but are too far south to be seen from Paris.




Below and left of the Sagittarius 'Teapot' is the glowing gas cloud **M8**, the '**Lagoon Nebula**'. It is a star-forming region where gas and dust have recently gathered into new stars. ('Recently' = the past million years or so.) Ultraviolet light from one particularly hot star is lighting up the leftover gas, making it glow. On colour photos it appears pink due to hydrogen atoms fluorescing in the UV light. Below M8 is M20, the Trifid Nebula, small glowing patch in binoculars, also a pink hydrogen region in photos. Other nearby nebulae (gas and dust clouds) are M16 and M17.



Globular clusters, spherical clusters of ancient stars, are found throughout the region. The brightest is **M4** by Antares. It is also one of the closest at 10 000 l.y. away. In binoculars and small telescopes 'globs' appear as round fuzzy spots. Others marked on the chart are M9, M10, M12, M14, M19, M22, M55, M54, M62, M80 and NGC 6541. The concentration of globular clusters in this area was an early clue that the centre of the galaxy lay in this direction.

This part of the Milky Way is broad and bright as we are looking to the centre of the galaxy. The actual centre, 27 000 light years away, is hidden from our view by intervening dust clouds. The nearer dust clouds make gaps and slots along the Milky Way. The hub of the galaxy is a great sphere of stars, called the 'central bulge'. Some of the central bulge is glimpsed in gaps between the dust clouds. At the very centre is a black hole four million times the sun's mass but only the size of our solar system. All big galaxies have a massive black hole at their centre.




Southern Evening Sky in September

The chart shows the southern and southwest sky. Interesting star clusters and nebulae are indicated with asterisks. They are described on the other side of this page.

Chart produced by Guide 8 software; www.projectpluto.com. Labels added by Alan Gilmore, University of Canterbury's Mt John Observatory
 P.O. Box 56, Lake Tekapo 7945, New Zealand. www.canterbury.ac.nz

Interesting Objects in the Southern Sky

Large & Small Clouds of Magellan (LMC & SMC) appear as two luminous patches, easily seen by eye in a dark sky. They are two galaxies like the Milky Way but much smaller. Each is made of billions of stars. The Large Cloud contains many clusters of young luminous stars seen as patches of light in binoculars and telescopes. The LMC is about 160 000 light years away and the SMC 200 000 l.y away, both very close for galaxies. (1 light year is about 10 000 billion km, 10^{13} km.)



47 Tucanae, looks like a faint fuzzy star on the edge of the SMC. It is a globular cluster, a ball of millions of stars. A telescope is needed to see a peppering of stars around the edge of the cluster. Though it appears on the edge of the SMC it is one-tenth the distance, 15 000 light years away, and is has no connection to the Small Cloud. Globular clusters are mostly very old, 10 billion years or more; at least twice the age of the sun. **Omega Centauri**, right of the Pointers, is a similar cluster around 17 000 light years away.




Tarantula nebula is a glowing gas cloud in the LMC. The gas glows in the ultra-violet light from a cluster of very hot stars at centre of the nebula. The cloud is about 800 light years across. It is easily seen in binoculars and can be seen by eye on moonless nights. This nebula is one of the brightest known. If it was as close as the Orion nebula (in The Pot's handle in the summer sky) then it would be as bright as the full moon.

Canopus is the second brightest star. It is 14 000 times brighter than the sun and 300 light years away. Sirius, in the eastern dawn sky, is the brightest star in the sky. The planets Venus and Jupiter are brighter.

Alpha Centauri, the brighter Pointer, is the closest naked-eye star, 4.3 light-years away. Alpha Centauri is a binary star: two stars about the same size as the sun orbiting around each other in 80 years. A telescope that magnifies 50x splits the pair. (A very faint and slightly closer star, Proxima Centauri, orbits a quarter of a light-year, or 15 000 Sun-earth distances, from the Alpha pair.)

Coalsack Nebula is a cloud of dust and gas about 600 light years away, dimming the more distant stars in the Milky Way. Many similar 'dark nebulae' can be seen, appearing as slots and holes in the Milky Way. These clouds of dust and gas eventually coalesce into clusters of stars.

The Jewel Box is a compact cluster of young luminous stars about 7000 light years away. The cluster formed less than 16 million years ago. It is best seen in a telescope. To the eye it looks like a faint star.



Eta Carinae nebula is a glowing gas cloud about 8000 light years away. The golden star in the cloud, visible in binoculars, is Eta Carinae. (Eta is the Greek 'e'.) It is estimated to be to be 80 times heavier than the sun and four million times brighter but is dimmed by dust clouds around it. It is expected to explode as a supernova any time in the next few thousand years. Many star clusters are found in this part of the sky.

The **Southern Pleiades** is a newish name for a cluster of stars at the right-hand point of the 'Diamond Cross'. It is formally the Theta Carinae cluster, after its brightest star but is also known as the 'Five of Diamonds' cluster, the reason obvious when it is seen in a telescope. It is much fainter and smaller than the real Pleiades in Taurus but a nice sight in binoculars. The cluster is about 500 light years away and is around 30 million years old.

NGC 2516, left of the False Cross, looks to the eye like a tailless comet. It is a nice sight in binoculars. The cluster is about 1200 light years away and 110 million years old.