

QUESTION ONE (8 marks)

There are three isomers of C_5H_{12} .

- (a) Draw the structure of each isomer.

--	--	--

- (b) One of the alkane isomers from part (a) reacts with Cl_2 in the presence of light to give a single monohalogenated organic product. Draw the structure of that product.


--

- (c) Another one of the isomers from part (a) reacts with Cl_2 in the presence of light to give four different monochloro alkanes. One of these is a secondary chloroalkane.

- (i) Draw the structure of this chloroalkane.

--

- (ii) Draw the structures of alkenes produced when the secondary chloroalkane from part (i) reacts with KOH in ethanol.



- (iii) Circle the alkene from part (ii) above that is formed in the greatest amount. Justify your answer.

QUESTION TWO (4 marks)

There are five C_5H_{10} constitutional (structural) isomers that are alkenes.

- (a) One of the five alkenes exists as *cis-trans* stereoisomers. Draw the structure of the *cis* isomer of this alkene.




- (b) Explain why the alkene in part (a) exists as *cis-trans* isomers whereas the isomeric alkene that has the same carbon skeleton does not.

- (c) Two C_5H_{10} alkenes have the same carbon skeleton, but neither exists as *cis-trans* isomers. One of these reacts with HCl to form a tertiary chloroalkane as the major product.

- (i) Draw the structure of this alkene;



- (ii) Draw the structure of the chloroalkane described above.



QUESTION THREE (12 marks)

This question is concerned with four alcohols and their physical and chemical properties. These alcohols have the formula $C_4H_{10}O$ and their boiling points are given in the table below.

Isomer	Boiling point	Reaction with $Cr_2O_7^{2-}$?
A	117 °C	Yes
B	102 °C	Yes
C	98 °C	Yes
D	82 °C	No

(a) (4 marks) Draw the structures of each of the four isomers.


(b) (2 marks) On the basis of your structures and the information given above, how does branching affect the attractive forces between the molecules? *Justify your answer.*

(c) (4 marks) Identify isomers **A** and **D** in your answer to part (a) above. *Justify your answer.*


(d) (2 marks) What additional information would you need to identify isomers **B** and **C**?

QUESTION FOUR (5 marks)

$[B_7H_7]^{2-}$ is a **pentagonal bipyramid** (shown below without the H atoms) with ten triangular faces. It has two types of B atoms; two axial (ax) and five equatorial (eq). A **nido-pentagonal bipyramid** is missing one of these vertices/atoms.




The cluster $[B_5NH_6]^{2-}$, in which one of the B atoms has been replaced by an N atom, is predicted to be a *nido*-pentagonal bipyramid. Sketch the possible isomers for this ion by writing B or N over the appropriate vertices in the polyhedra given below. You may not need to use all of the polyhedra to show all of the isomers.




QUESTION FIVE (12 marks)

(a) Draw ONE Lewis structure and the 3-dimensional molecular shape for each of the following molecules:

(i) Hydrogen peroxide (H_2O_2);



(ii) Trioxidane (H_2O_3);



(b) (i) Draw ONE Lewis structure and the 3-dimensional molecular shape for hyponitrous acid (a symmetric molecule with formula $\text{N}_2(\text{OH})_2$);



(ii) Hyponitrous acid can be synthesised as both the *cis* and *trans* isomers, but the *trans* isomer is the most stable. Give a reason in terms of electron pair repulsion theory to explain this.

- (c) (i) Draw TWO Lewis structures for nitrogen dioxide (NO_2) and the 3-dimensional molecular shape for one of these.

--	--	--

- (ii) The "formal charge" is the number of valence electrons in the atom, minus the number of lone pair electrons at that atom in the Lewis structure, minus the number of bonds to the atom in the Lewis structure. Formal charge can be used to help explain where electrons are likely to be found on atoms in a molecule. Identify your structure in part (i) that formal charge suggests is most likely.
- (iii) The O-N-O bond angle is actually 134.3° . Does this value support your proposed Lewis structure? Justify your answer.

QUESTION SIX (10 marks)

Silver nitrate is used in volumetric analysis to determine the concentration of chloride ions in an aqueous solution. Because of the high cost of AgNO_3 , a student uses an available supply of $0.0500 \text{ mol L}^{-1}$ AgNO_3 solution and some solid AgNO_3 to prepare 100.0 mL of $0.0750 \text{ mol L}^{-1}$ AgNO_3 . She prepares the solution by:

- (i) pipetting exactly 50.00 mL of the $0.0500 \text{ mol L}^{-1}$ AgNO_3 solution into a 100.0 mL volumetric flask;
 - (ii) adding an appropriate mass of AgNO_3 ;
 - (iii) diluting the solution to exactly 100.0 mL .
- (a) What mass of AgNO_3 should be added in step (ii)? [$M(\text{AgNO}_3) = 169.9 \text{ g mol}^{-1}$]

- (b) Solid MgCl_2 (0.100 g) was then added to the solution. Assuming no change in the total volume, what is the concentration of each of the following species?
[$M(\text{MgCl}_2) = 95.2 \text{ g mol}^{-1}$]



QUESTION SEVEN (7 marks)

The Kjeldahl method can be used to determine the percentage of nitrogen in meat and other organic products. A 0.0986 g sample was heated with concentrated sulfuric acid for two hours to oxidise organic matter and convert all nitrogen to ammonium ions. The solution was then made strongly basic by adding excess sodium hydroxide solution producing ammonia. The ammonia was then distilled into 50.00 mL of 0.1010 mol L⁻¹ HNO₃. Exactly 23.45 mL of 0.1500 mol L⁻¹ NaOH was required to neutralise the excess acid.

Calculate the amount (moles) of NH₃ that was distilled into the HNO₃ and hence determine the percentage of N in the original sample.

QUESTION EIGHT (6 marks)

For each of the following compounds, state with brief explanation whether its solubility in water will increase, decrease or be unaffected by a decrease in pH:

(a) PbSO_4

(b) AgCl

(c) CuS

QUESTION NINE (5 marks)

A compound consists of 14.29% carbon, 57.14% oxygen, 1.19% hydrogen and an element X having the same number of moles as there are moles of carbon.

(a) Identify X.



(b) Determine the empirical formula of the compound.

(c) Suggest the likely identity of the compound.

QUESTION ELEVEN (13 marks)

When aqueous ammonia is added drop-wise to a copper sulfate solution, a blue solid of copper hydroxide forms. As further ammonia is added, the solid redissolves and a dark blue solution forms.

The following equilibria explain these observations:



- (a) Give the equilibrium expression for each of these processes.
- (b) Explain why copper hydroxide can form when ammonia is added.
- (c) The equilibrium constant for process A is 2.20×10^{-20} while the equilibrium constant for process B is 1.2×10^{13} . Use these values to explain why initially a precipitate forms with limited ammonia but, when excess ammonia is added, the solid redissolves to form the dark blue ammonia complex.
- (d) What happens to the concentration of each of the following species, once equilibrium is re-established, upon addition of $\text{Cu}^{2+}_{(\text{aq})}$ to the dark blue solution?
- (i) $[\text{Cu}(\text{NH}_3)_4]^{2+}_{(\text{aq})}$
- (ii) NH_3
- (iii) $\text{Cu}^{2+}_{(\text{aq})}$

QUESTION TWELVE (6 marks)

The average chemical formula for common diesel fuel is $C_{12}H_{26}$. Dodecane ($C_{12}H_{26}$) has an enthalpy of combustion of $-8072 \text{ kJ mol}^{-1}$ and a density of 0.745 g mL^{-1} . The enthalpy of combustion for a given substance is defined as the enthalpy change for the reaction of one mole of the substance with oxygen to form $CO_2(g)$ and $H_2O(l)$. $M(C_{12}H_{26}) = 170 \text{ g mol}^{-1}$.

- (a) (2 marks) Write down a balanced equation for the combustion of dodecane.
- (b) (2 marks) Calculate the energy density, expressed as kJ of heat given off in combustion per litre of fuel (kJ/L) for dodecane.
- (c) (2 marks) What mass of CO_2 is produced in order to generate 15,000 kJ of energy?

PERIODIC TABLE OF THE ELEMENTS

																		18											
																		1 H 1.0	2										
																		Atomic Number		Molar Mass / g mol ⁻¹				13	14	15	16	17	18
3	4													5	6	7	8	9	10										
Li 6.9	Be 9.0													B 10.8	C 12.0	N 14.0	O 16.0	F 19.0	Ne 20.2										
11	12													13	14	15	16	17	18										
Na 23.0	Mg 24.3													Al 27.0	Si 28.1	P 31.0	S 32.1	Cl 35.5	Ar 40.0										
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36												
K 39.1	Ca 40.1	Sc 45.0	Ti 47.9	V 50.9	Cr 52.0	Mn 54.9	Fe 55.9	Co 58.9	Ni 58.7	Cu 63.5	Zn 65.4	Ga 69.	Ge 72.6	As 74.9	Se 79.0	Br 79.9	Kr 83.8												
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54												
Rb 85.5	Sr 87.6	Y 88.9	Zr 91.2	Nb 92.9	Mo 95.9	Tc 98.9	Ru 101	Rh 103	Pd 106	Ag 108	Cd 112	In 115	Sn 119	Sb 122	Te 128	I 127	Xe 131												
55	56	57-71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86												
Cs 133	Ba 137	Lanthanide Series	Hf 179	Ta 181	W 184	Re 186	Os 190	Ir 192	Pt 195	Au 197	Hg 201	Tl 204	Pb 207	Bi 209	Po 210	At 210	Rn 222												
87	88	89-103	104	105	106	107	108	109																					
Fr 223	Ra 226	Actinide Series	Rf 261	Db 262	Sg 263	Bh 262	Hs 265	Mt 266																					

	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
Lanthanide Series	La 139	Ce 140	Pr 141	Nd 144	Pm 145	Sm 150	Eu 152	Gd 157	Tb 159	Dy 163	Ho 165	Er 167	Tm 169	Yb 173	Lu 175
Actinide Series	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
	Ac 227	Th 232	Pa 231	U 238	Np 237	Pu 244	Am 243	Cm 247	Bk 247	Cf 251	Es 252	Fm 257	Md 258	No 255	Lr 262